

Question 6

1) The Maclaurin series for e^x is $e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots + \frac{x^n}{n!} + \dots$. The continuous function f is defined by $f(x) = \frac{e^{(x-1)^2} - 1}{(x-1)^2}$ for $x \neq 1$ and $f(1) = 1$. The function f has derivatives of all orders at $x = 1$.

- Write the first four nonzero terms and the general term of the Taylor series for $e^{(x-1)^2}$ about $x = 1$.
- Use the Taylor series found in part (a) to write the first four nonzero terms and the general term of the Taylor series for f about $x = 1$.
- Use the ratio test to find the interval of convergence for the Taylor series found in part (b).
- Use the Taylor series for f about $x = 1$ to determine whether the graph of f has any points of inflection.

Question 6

2) A function f has derivatives of all orders at $x = 0$. Let $P_n(x)$ denote the n th-degree Taylor polynomial for f about $x = 0$.

- It is known that $f(0) = -4$ and that $P_1\left(\frac{1}{2}\right) = -3$. Show that $f'(0) = 2$.
- It is known that $f''(0) = -\frac{2}{3}$ and $f'''(0) = \frac{1}{3}$. Find $P_3(x)$.
- The function h has first derivative given by $h'(x) = f(2x)$. It is known that $h(0) = 7$. Find the third-degree Taylor polynomial for h about $x = 0$.

Question 6

3) The Maclaurin series for a function f is given by $\sum_{n=1}^{\infty} \frac{(-3)^{n-1}}{n} x^n = x - \frac{3}{2}x^2 + 3x^3 - \dots + \frac{(-3)^{n-1}}{n} x^n + \dots$ and converges to $f(x)$ for $|x| < R$, where R is the radius of convergence of the Maclaurin series.

- Use the ratio test to find R .
- Write the first four nonzero terms of the Maclaurin series for f' , the derivative of f . Express f' as a rational function for $|x| < R$.
- Write the first four nonzero terms of the Maclaurin series for e^x . Use the Maclaurin series for e^x to write the third-degree Taylor polynomial for $g(x) = e^x f(x)$ about $x = 0$.

4)

Question 6

The Maclaurin series for the function f is given by $f(x) = \sum_{n=2}^{\infty} \frac{(-1)^n (2x)^n}{n-1}$ on its interval of convergence.

- (a) Find the interval of convergence for the Maclaurin series of f . Justify your answer.
- (b) Show that $y = f(x)$ is a solution to the differential equation $xy' - y = \frac{4x^2}{1+2x}$ for $|x| < R$, where R is the radius of convergence from part (a).

5)

Question 6

Let f be a function with derivatives of all orders and for which $f(2) = 7$. When n is odd, the n th derivative of f at $x = 2$ is 0. When n is even and $n \geq 2$, the n th derivative of f at $x = 2$ is given by $f^{(n)}(2) = \frac{(n-1)!}{3^n}$.

- (a) Write the sixth-degree Taylor polynomial for f about $x = 2$.
- (b) In the Taylor series for f about $x = 2$, what is the coefficient of $(x-2)^{2n}$ for $n \geq 1$?
- (c) Find the interval of convergence of the Taylor series for f about $x = 2$. Show the work that leads to your answer.

6)

The function f is defined by the power series

$$f(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n+1)!} = 1 - \frac{x^2}{3!} + \frac{x^4}{5!} - \frac{x^6}{7!} + \cdots + \frac{(-1)^n x^{2n}}{(2n+1)!} + \cdots$$

for all real numbers x .

- (a) Find $f'(0)$ and $f''(0)$. Determine whether f has a local maximum, a local minimum, or neither at $x = 0$. Give a reason for your answer.
- (b) Show that $1 - \frac{1}{3!}$ approximates $f(1)$ with error less than $\frac{1}{100}$.
- (c) Show that $y = f(x)$ is a solution to the differential equation $xy' + y = \cos x$.