

| | | | | | |
|--------------------|-----|---|---|----|----|
| t (years) | 2 | 3 | 5 | 7 | 10 |
| $H(t)$ (meters) | 1.5 | 2 | 6 | 11 | 15 |

4. The height of a tree at time t is given by a twice-differentiable function H , where $H(t)$ is measured in meters and t is measured in years. Selected values of $H(t)$ are given in the table above.
- Use the data in the table to estimate $H'(6)$. Using correct units, interpret the meaning of $H'(6)$ in the context of the problem.
 - Explain why there must be at least one time t , for $2 < t < 10$, such that $H'(t) = 2$.
 - Use a trapezoidal sum with the four subintervals indicated by the data in the table to approximate the average height of the tree over the time interval $2 \leq t \leq 10$.

| | | | | |
|-------------------------|------|------|-----|-----|
| h (feet) | 0 | 2 | 5 | 10 |
| $A(h)$ (square feet) | 50.3 | 14.4 | 6.5 | 2.9 |

1. A tank has a height of 10 feet. The area of the horizontal cross section of the tank at height h feet is given by the function A , where $A(h)$ is measured in square feet. The function A is continuous and decreases as h increases. Selected values for $A(h)$ are given in the table above.
- Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the volume of the tank. Indicate units of measure.
 - Does the approximation in part (a) overestimate or underestimate the volume of the tank? Explain your reasoning.

| | | | | | |
|---------------------------|------|------|-----|-----|-----|
| t (hours) | 0 | 1 | 3 | 6 | 8 |
| $R(t)$ (liters / hour) | 1340 | 1190 | 950 | 740 | 700 |

1. Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.
- Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
 - Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
 - Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
 - For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.

| | | | | | | | |
|--------------------|---|-----|-----|------|------|------|------|
| t (minutes) | 0 | 1 | 2 | 3 | 4 | 5 | 6 |
| $C(t)$ (ounces) | 0 | 5.3 | 8.8 | 11.2 | 12.8 | 13.8 | 14.5 |

3. Hot water is dripping through a coffeemaker, filling a large cup with coffee. The amount of coffee in the cup at time t , $0 \leq t \leq 6$, is given by a differentiable function C , where t is measured in minutes. Selected values of $C(t)$, measured in ounces, are given in the table above.
- Use the data in the table to approximate $C'(3.5)$. Show the computations that lead to your answer, and indicate units of measure.
 - Is there a time t , $2 \leq t \leq 4$, at which $C'(t) = 2$? Justify your answer.
 - Use a midpoint sum with three subintervals of equal length indicated by the data in the table to approximate the value of $\frac{1}{6} \int_0^6 C(t) dt$. Using correct units, explain the meaning of $\frac{1}{6} \int_0^6 C(t) dt$ in the context of the problem.

| | | | | | |
|-----------------------------|------|------|------|------|------|
| t (minutes) | 0 | 4 | 9 | 15 | 20 |
| $W(t)$ (degrees Fahrenheit) | 55.0 | 57.1 | 61.8 | 67.9 | 71.0 |

- 5
1. The temperature of water in a tub at time t is modeled by a strictly increasing, twice-differentiable function W , where $W(t)$ is measured in degrees Fahrenheit and t is measured in minutes. At time $t = 0$, the temperature of the water is 55°F . The water is heated for 30 minutes, beginning at time $t = 0$. Values of $W(t)$ at selected times t for the first 20 minutes are given in the table above.
- (a) Use the data in the table to estimate $W'(12)$. Show the computations that lead to your answer. Using correct units, interpret the meaning of your answer in the context of this problem.
- (b) Use the data in the table to evaluate $\int_0^{20} W'(t) dt$. Using correct units, interpret the meaning of $\int_0^{20} W'(t) dt$ in the context of this problem.
- (c) For $0 \leq t \leq 20$, the average temperature of the water in the tub is $\frac{1}{20} \int_0^{20} W(t) dt$. Use a left Riemann sum with the four subintervals indicated by the data in the table to approximate $\frac{1}{20} \int_0^{20} W(t) dt$. Does this approximation overestimate or underestimate the average temperature of the water over these 20 minutes? Explain your reasoning.

| | | | | | |
|-----------------------------|----|----|----|----|----|
| t (minutes) | 0 | 2 | 5 | 9 | 10 |
| $H(t)$ (degrees Celsius) | 66 | 60 | 52 | 44 | 43 |

- 6
2. As a pot of tea cools, the temperature of the tea is modeled by a differentiable function H for $0 \leq t \leq 10$, where time t is measured in minutes and temperature $H(t)$ is measured in degrees Celsius. Values of $H(t)$ at selected values of time t are shown in the table above.
- (a) Use the data in the table to approximate the rate at which the temperature of the tea is changing at time $t = 3.5$. Show the computations that lead to your answer.
- (b) Using correct units, explain the meaning of $\frac{1}{10} \int_0^{10} H(t) dt$ in the context of this problem. Use a trapezoidal sum with the four subintervals indicated by the table to estimate $\frac{1}{10} \int_0^{10} H(t) dt$.

| | | | | | |
|---------------------------------|---|---|----|----|----|
| t (hours) | 0 | 2 | 5 | 7 | 8 |
| $E(t)$ (hundreds of entries) | 0 | 4 | 13 | 21 | 23 |

7. 2. A zoo sponsored a one-day contest to name a new baby elephant. Zoo visitors deposited entries in a special box between noon ($t = 0$) and 8 P.M. ($t = 8$). The number of entries in the box t hours after noon is modeled by a differentiable function E for $0 \leq t \leq 8$. Values of $E(t)$, in hundreds of entries, at various times t are shown in the table above.

(a) Use the data in the table to approximate the rate, in hundreds of entries per hour, at which entries were being deposited at time $t = 6$. Show the computations that lead to your answer.

(b) Use a trapezoidal sum with the four subintervals given by the table to approximate the value of $\frac{1}{8} \int_0^8 E(t) dt$.

Using correct units, explain the meaning of $\frac{1}{8} \int_0^8 E(t) dt$ in terms of the number of entries.

| | | | | | |
|--------|---|---|----|---|----|
| x | 2 | 3 | 5 | 8 | 13 |
| $f(x)$ | 1 | 4 | -2 | 3 | 6 |

8. 5. Let f be a function that is twice differentiable for all real numbers. The table above gives values of f for selected points in the closed interval $2 \leq x \leq 13$.

(a) Estimate $f'(4)$. Show the work that leads to your answer.

(b) Evaluate $\int_2^{13} (3 - 5f'(x)) dx$. Show the work that leads to your answer.

(c) Use a left Riemann sum with subintervals indicated by the data in the table to approximate $\int_2^{13} f(x) dx$.

Show the work that leads to your answer.

| | | | | | | | |
|-----------------|-----|-----|-----|-----|-----|----|---|
| t (hours) | 0 | 1 | 3 | 4 | 7 | 8 | 9 |
| $L(t)$ (people) | 120 | 156 | 176 | 126 | 150 | 80 | 0 |

2. Concert tickets went on sale at noon ($t = 0$) and were sold out within 9 hours. The number of people waiting in line to purchase tickets at time t is modeled by a twice-differentiable function L for $0 \leq t \leq 9$. Values of $L(t)$ at various times t are shown in the table above.
- (a) Use the data in the table to estimate the rate at which the number of people waiting in line was changing at 5:30 P.M. ($t = 5.5$). Show the computations that lead to your answer. Indicate units of measure.
- (b) Use a trapezoidal sum with three subintervals to estimate the average number of people waiting in line during the first 4 hours that tickets were on sale.
- (c) For $0 \leq t \leq 9$, what is the fewest number of times at which $L'(t)$ must equal 0? Give a reason for your answer.

| | | | | | | |
|------------------------------|-----|-----|-----|-----|-----|-----|
| t (minutes) | 0 | 2 | 5 | 7 | 11 | 12 |
| $r'(t)$ (feet per minute) | 5.7 | 4.0 | 2.0 | 1.2 | 0.6 | 0.5 |

5. The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function r of time t , where t is measured in minutes. For $0 < t < 12$, the graph of r is concave down. The table above gives selected values of the rate of change, $r'(t)$, of the radius of the balloon over the time interval $0 \leq t \leq 12$. The radius of the balloon is 30 feet when $t = 5$. (Note: The volume of a sphere of radius r is given by $V = \frac{4}{3}\pi r^3$.)
- (a) Estimate the radius of the balloon when $t = 5.4$ using the tangent line approximation at $t = 5$. Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when $t = 5$. Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate $\int_0^{12} r'(t) dt$. Using correct units, explain the meaning of $\int_0^{12} r'(t) dt$ in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than $\int_0^{12} r'(t) dt$? Give a reason for your answer.